

# Bcl-2-mediated alterations in endoplasmic reticulum $\text{Ca}^{2+}$ analyzed with an improved genetically encoded fluorescent sensor

Amy E. Palmer\*, Can Jin<sup>†</sup>, John C. Reed<sup>†</sup>, and Roger Y. Tsien\*<sup>‡</sup>

\*Department of Pharmacology and Howard Hughes Medical Institute, University of California at San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0647; and <sup>†</sup>The Burnham Institute, La Jolla, CA 92037

Contributed by Roger Y. Tsien, November 1, 2004

The endoplasmic reticulum (ER) serves as a cellular storehouse for  $\text{Ca}^{2+}$ , and  $\text{Ca}^{2+}$  released from the ER plays a role in a host of critical signaling reactions, including exocytosis, contraction, metabolism, regulation of transcription, fertilization, and apoptosis. Given the central role played by the ER, our understanding of these signaling processes could be greatly enhanced by the ability to image  $[\text{Ca}^{2+}]_{\text{ER}}$  directly in individual cells. We created a genetically encoded  $\text{Ca}^{2+}$  indicator by redesigning the binding interface of calmodulin and a calmodulin-binding peptide. The sensor has improved reaction kinetics and a  $K_d$  ideal for imaging  $\text{Ca}^{2+}$  in the ER and is no longer perturbed by large excesses of native calmodulin. Importantly, it provides a significant improvement over all previous methods for monitoring  $[\text{Ca}^{2+}]_{\text{ER}}$  and has been used to directly show that, in MCF-7 breast cancer cells, the antiapoptotic protein B cell lymphoma 2 (Bcl-2) (*i*) lowers  $[\text{Ca}^{2+}]_{\text{ER}}$  by increasing  $\text{Ca}^{2+}$  leakage under resting conditions and (*ii*) alters  $\text{Ca}^{2+}$  oscillations induced by ATP, and that acute inhibition of Bcl-2 by the green tea compound epigallocatechin gallate results in an increase in  $[\text{Ca}^{2+}]_{\text{ER}}$  due to inhibition of Bcl-2-mediated  $\text{Ca}^{2+}$  leakage.

calcium | cameleon

To date, researchers have attempted to use a number of strategies to deduce the role of  $[\text{Ca}^{2+}]_{\text{ER}}$  in signaling processes, but all possess shortcomings (1, 2). Conventional methods use cytosolic small-molecule  $\text{Ca}^{2+}$  indicators to infer that an observed signal likely originated from the endoplasmic reticulum (ER). However, as more studies identify the complex interplay among levels of  $\text{Ca}^{2+}$  in the ER and  $\text{Ca}^{2+}$  influx (3), as well as a close interaction network between the ER and mitochondria (4, 5), it seems increasingly important for our understanding of signaling processes that organelle  $\text{Ca}^{2+}$  be observed directly to differentiate  $\text{Ca}^{2+}$  signals from different organelles.  $\text{Ca}^{2+}$  indicator dyes can be loaded at 37°C rather than 25°C, thus favoring the loading of internal compartments. However, this method loads all internal compartments (not just the ER), and the remaining cytosolic dye must be effectively removed, or quenched. ER-targeted aequorin probes (6) can be genetically targeted to the ER to measure  $[\text{Ca}^{2+}]_{\text{ER}}$  directly, but the aequorin probes are luminescent and therefore not bright enough for single-cell imaging without specialized equipment. Perhaps more problematic is that for aequorin to be active it must be reconstituted with the coelenterazine cofactor, and to do so the ER must initially be completely depleted of  $\text{Ca}^{2+}$ . Finally, the low-affinity FRET-based cameleon (7) YC4.3ER can also be genetically targeted and enables single-cell imaging, but its  $K_d$  for  $\text{Ca}^{2+}$  is not well suited to monitor changes in  $[\text{Ca}^{2+}]_{\text{ER}}$ , and therefore it has a low sensitivity.

Given the critical role of apoptosis in human diseases as well as normal physiology and the increasing evidence that the balance of B cell lymphoma 2 (Bcl-2) family proteins plays a role in the tumorigenesis of a number of cancers (8), a fundamental need exists to better understand how these proteins interact and the multiple mechanisms by which they influence apoptotic

cascades. An important aspect of this is identifying inhibitors of different Bcl-2 family members that could serve as tools to perturb and thereby better our understanding of the signaling cascades, as well as to act as potential therapeutic agents. We set out to generate an indicator for  $[\text{Ca}^{2+}]_{\text{ER}}$  so that we could examine the role of  $[\text{Ca}^{2+}]_{\text{ER}}$  in the regulation of Bcl-2 family-mediated apoptosis. Although most studies have focused on the role of the mitochondria in apoptosis, recently it has been shown that Bcl-2 family members can localize to the ER and alter  $[\text{Ca}^{2+}]_{\text{ER}}$ , thus adding another dimension to the role of Bcl-2 family members in apoptosis (9–13). Importantly, it has been shown that changes in  $[\text{Ca}^{2+}]_{\text{ER}}$  can directly influence the propensity of a given stimulus to lead to apoptosis (12, 14), although there have been a number of conflicting reports (15) as to exactly how the antiapoptotic protein Bcl-2 influences  $[\text{Ca}^{2+}]_{\text{ER}}$ . Of particular interest is a recent report that Bcl-2 functionally interacts with inositol 1,4,5-trisphosphate ( $\text{IP}_3$ ) receptors ( $\text{IP}_3\text{Rs}$ ) in WEHI7.2 T cells, suggesting that this interaction might be the mechanism by which Bcl-2 alters ER  $\text{Ca}^{2+}$  homeostasis (16). The creation of a robust and reliable sensor to directly monitor  $[\text{Ca}^{2+}]_{\text{ER}}$  could help alleviate some of the uncertainties associated with indirect  $\text{Ca}^{2+}$  measurements and would provide a significant advantage in probing questions related to  $\text{Ca}^{2+}$  homeostasis in the ER.

## Materials and Methods

Peptide synthesis and Biacore (Neuchâtel, Switzerland) experiments were performed as described in *Supporting Text*, which is published as supporting information on the PNAS web site. The mutant peptide and calmodulins (CaMs) were cloned between a truncated enhanced cyan fluorescent protein (CFP) and citrine fluorescent protein (17), as described (7). The constructs were cloned between the *Bam*HI/*Eco*RI sites in pRSETB or pBAD for protein purification and pcDNA3 for expression in mammalian cells. To generate an ER-targeted cameleon, the calreticulin signal sequence MLLPVLLGLLGAAD was added 5' to CFP, and an ER retention sequence, KDEL, was added to the 3' end of citrine.

For protein purification, cameleons were expressed in either JM109 (Stratagene) or LMG194 (Invitrogen) and grown overnight at 25°C. Constructs in pBAD/LMG194 were induced with 0.2% arabinose. Protein was extracted with Bacterial Protein Extraction Reagent (Pierce), purified via an N-terminal 6×His-

Freely available online through the PNAS open access option.

Abbreviations: ER, endoplasmic reticulum; SERCA, sarcoendoplasmic reticulum  $\text{Ca}^{2+}$  ATPase; CaM, calmodulin; skMLCK, skeletal muscle myosin light chain kinase; Bcl-2, B cell lymphoma 2;  $\text{IP}_3$ , inositol 1,4,5-trisphosphate;  $\text{IP}_3\text{R}$ ,  $\text{IP}_3$  receptor; EGCG, epigallocatechin gallate; EC, epicatechin; CFP, cyan fluorescent protein; HBSS, Hanks' balanced salt solution; D1, Design 1.

Data deposition: The sequence reported in this paper has been deposited in the GenBank database (accession no. AY796115).

<sup>‡</sup>To whom correspondence should be addressed. E-mail: rtsien@ucsd.edu.

© 2004 by The National Academy of Sciences of the USA


tag by using Ni-NTA agarose, and buffer-exchanged into 10 mM 4-morpholinepropanesulfonic acid/100 mM NaCl, pH 7.4, by using Amicon Centricon-30 columns (Millipore). Absorbance measurements were conducted on a Cary 3E UV-Visible spectrometer (Varian). Fluorescence measurements were conducted on a Spex (Spex Industries, Metuchen, NJ) Fluorolog-3 Fluorimeter (Horiba group) at a cameleon concentration of 0.4  $\mu\text{M}$ .  $\text{Ca}^{2+}$ /EGTA, and  $\text{Ca}^{2+}$ /*N*-(2-hydroxyethyl)ethylene dinitrilo-*N,N,N'*-triacetic acid (HEEDTA) buffers were prepared as described (18, 19) and were used in  $\text{Ca}^{2+}$  titrations to achieve concentrations of  $<50 \mu\text{M}$ . All solutions of  $>50 \mu\text{M}$  were unbuffered.

Stopped-flow experiments were performed on an Applied Photophysics (Surrey, U.K.) Stopped Flow. Equal volumes of cameleon (final concentration of 0.8  $\mu\text{M}$ ) and  $\text{Ca}^{2+}$  (various concentrations) were rapidly mixed, and the fluorescence was monitored by using an excitation wavelength of 420 nm and an emission cutoff filter of 500 nm. The observed first-order rate constant ( $k_{\text{obs}}$ ) was calculated from each averaged ( $n > 4$ ) data set by nonlinear regression analysis.

Cell-culturing details are provided in *Supporting Text*. For imaging, cells were rinsed twice and then maintained in Hanks' balanced salt solution (HBSS) with 20 mM Hepes and 2 g/liter D-glucose at pH 7.4.  $\text{Ca}^{2+}$ -free solutions were generated by adding 0.5 mM EGTA, 1 mM  $\text{Mg}^{2+}$ , 1 g/liter glucose, and 20 mM Hepes to  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ -free HBSS (GIBCO). For experiments in which both the ER and cytosolic  $\text{Ca}^{2+}$  levels were monitored, cells were treated with 4  $\mu\text{M}$  fura-2-AM (Molecular Probes) with 0.04% Pluronic F-127 in HBSS for 30 min at room temperature. Cells were then washed with HBSS and allowed to incubate at room temperature for 15 min to ensure cleavage of the acetoxyethyl ester. (–)Epigallocatechin gallate (EGCG) and (–)epicatechin (EC) were purchased from Sigma, dissolved in 100% DMSO, and stored at  $-80^\circ\text{C}$  until use.

To calibrate the cameleon probe in the ER,  $R_{\text{min}}$  was obtained by treating the cells with 3 mM EGTA and 2  $\mu\text{M}$  ionomycin, and  $R_{\text{max}}$  was determined by treating cells with 25  $\mu\text{M}$  digitonin, followed by 5–10 mM  $\text{Ca}^{2+}$ , 1 mM ATP, and 1 mM  $\text{Mg}^{2+}$ . The  $R_{\text{min}}$  and  $R_{\text{max}}$  values were used to convert cell data to % FRET (of maximum). Finally, the apparent  $K_{d1}$  (0.58),  $K_{d2}$  (56.46),  $R_{\text{max}1}$  (0.28),  $R_{\text{max}2}$  (0.72),  $n_1$  (1.18), and  $n_2$  (1.67) values obtained from fitting the *in vitro* data with  $R = \{R_{\text{max}1}[\text{Ca}^{2+}]^{n_1}/(K_{d1}^{n_1} + [\text{Ca}^{2+}]^{n_1})\} + \{R_{\text{max}2}[\text{Ca}^{2+}]^{n_2}/(K_{d2}^{n_2} + [\text{Ca}^{2+}]^{n_2})\}$  were used in the calibration. To calibrate fura-2, cells were treated with 8  $\mu\text{M}$  ionomycin and 10 mM EGTA in  $\text{Ca}^{2+}$ -free HBSS to obtain  $R_{\text{min}}$ , followed by 2  $\mu\text{M}$  ionomycin and 20 mM  $\text{Ca}^{2+}$  in HBSS to obtain  $R_{\text{max}}$ . The *in situ*  $R_{\text{min}}$  and  $R_{\text{max}}$  values were adjusted (multiplied by 0.85) to account for the minimum viscosity effect (20). The standard equation:  $K_d[(R - R_{\text{min}})/(R_{\text{max}} - R)] \cdot S_f/S_b$  was used to convert the Fura-2 350:380 ratio to  $[\text{Ca}^{2+}]_{\text{cyt}}$ , where  $S_f$  and  $S_b$  are the emission intensity at 380 nm for  $\text{Ca}^{2+}$ -free and  $\text{Ca}^{2+}$ -bound fura-2, respectively.

Cells were imaged on a Zeiss Axiovert 200M microscope with a cooled charge-coupled device camera (Roper Scientific, Trenton, NJ), controlled by METAFLUOR 6.1 software (Universal Imaging, Downingtown, PA). Emission ratio imaging of the cameleon was accomplished by using a 436DF20 excitation filter, 450-nm dichroic mirror, and two emission filters (475/40 for enhanced CFP and 535/25 for citrine) controlled by a Lambda 10-2 filter changer (Sutter Instruments, Novato, CA). Excitation ratio imaging for fura-2 was accomplished by using 350/10 and 380/10 excitation filters, a 450-nm dichroic mirror, and a 535/45 emission filter. Fluorescence images were background corrected. Exposure times were typically 100–1,000 ms, and images were collected every 8–20 s.




**Fig. 1.** *In vitro* characterization of the mutant CaM and peptide and the resulting redesigned cameleon. (a) Biacore sensorgram of WT CaM binding to skMLCK (solid line) and four-charge reversal skMLCK (dashed line). CaM was injected over the surface in the presence of saturating  $\text{Ca}^{2+}$  starting at time  $t = 0$  for 360 s. (b) Binding of WT CaM (squares, dashed line) and mutant CaM (circles, solid line) to mutant skMLCK with four-charge reversals. The binding of CaM reached a steady state during the association phase, and therefore a Scatchard analysis was used to determine the dissociation constants (Fig. 8). (c) *In vitro* calcium titration curves of YC2.1 (diamonds, dashed line), YC3.3 (squares, dashed line), YC4.3 (triangles, dashed line), and D1 (circles, solid line), along with corresponding fits of the data. The gray box represents the typical range of  $[\text{Ca}^{2+}]$  in the ER, from resting to the depleted state. (d) Percent of the maximum FRET response of YC3.3 (squares) and D1 (circles) with increasing concentrations of WT CaM.

## Results

To create a sensor that would be appropriate for monitoring  $[\text{Ca}^{2+}]_{\text{ER}}$  in single living cells, we started with the original cameleon construct, comprised of two fluorescent proteins (CFP and citrine) and two sensing proteins [CaM and a CaM-binding peptide derived from skeletal muscle myosin light chain kinase (skMLCK)] that undergo a conformational change upon binding (7). Our goal was to redesign the binding interface between CaM and the peptide to generate highly specific protein/peptide pairs that would display a range of  $\text{Ca}^{2+}$  affinities and that would not be perturbed by endogenous proteins, such as WT CaM. This was particularly important, given that previous cameleons have suffered from perturbation by endogenous proteins (21, 22). Toward this end, we initially generated peptides that would not bind to WT CaM and then sought to reengineer CaM to reconstitute binding to the mutant peptides.

As a starting point for redesign, we targeted the six possible salt-bridge interactions identified in the NMR solution structure between CaM and the skMLCK peptide (23). A series of biotinylated peptides was synthesized in which a basic residue (either K or R) at the N and C termini of the peptide was replaced with an acidic E residue. Charge reversals were made sequentially, generating peptides with two, four, and six charge



**Fig. 2.** Comparison of the response of ER-targeted YC4.3 and D1 in HeLa cells. (a) Fluorescence image of D1ER in HeLa cells showing effective ER localization. (b) Emission ratio of D1 (red) and YC4.3 (blue) in the ER upon treatment with thapsigargin. (c) Emission ratio of D1 (red lines, representing two different cells) and YC4.3 (blue) in the ER upon treatment with ATP. (d) Simultaneous imaging of  $\text{Ca}^{2+}$  in the cytosol (using fura-2, green) and ER (D1, red). The left axis represents the excitation ratio of fura-2 (350/380 nm), and the right axis represents the emission ratio of the cameleon (citrine/CFP). (e) Same experimental conditions as in d but for an extended amount of time. In the continued presence of  $10 \mu\text{M}$  ATP, the ER begins to refill with  $\text{Ca}^{2+}$  after  $\approx 1,000$  s.

reversals until all of the salt-bridge interactions were destroyed. Binding of CaM to the peptides was assayed by using surface plasmon resonance (Biacore 3000), in which binding events are optically detected as a change in the refractive index at the surface. To accomplish this, biotinylated peptides were coupled to a streptavidin-coated chip, and CaM was injected over the surface in the presence of  $\text{Ca}^{2+}$  (Fig. 6, which is published as supporting information on the PNAS web site). Initially, we verified that binding of CaM to the WT skMLCK was  $\approx 10$  nM, comparable to previously reported values obtained in solution (Fig. 7, which is published as supporting information on the PNAS web site). The mutant peptide with four charge reversals displayed a dramatically lower affinity for CaM than the WT skMLCK peptide (Fig. 1a). Injection of increasing concentrations of CaM enabled determination of the  $K_d > 700 \mu\text{M}$  (Fig. 8, which is published as supporting information on the PNAS web site). Thus, four charge reversals were sufficient to decrease the affinity for WT CaM by  $>10^4$ .

A small library of CaM mutants in which the four complementary positions in CaM (E11, E84, E87, and E127) were changed to E, K, or R was then assayed for binding to both WT skMLCK and the mutant peptide. The best hit had a  $K_d$  of  $270 \mu\text{M}$ , and therefore a higher affinity than WT CaM for the mutant peptide (Fig. 1b). Upon sequencing, this mutant was identified as E11K, E84R, E87K, or E127E. Mutation of E127 to R did not increase the affinity of CaM for the peptide (data not shown).


The mutant CaM/peptide pair, designated as Design 1 (D1), was cloned between CFP and citrine to yield a reengineered cameleon. Under saturating conditions of  $\text{Ca}^{2+}$ , the cameleon showed a decrease in CFP emission and an increase in citrine emission, indicative of increased FRET. The magnitude of the FRET change was comparable to that observed for the original cameleons. Fig. 1c shows the  $\text{Ca}^{2+}$  titration curves of the design along with previous cameleons (YC2, YC3, and YC4). D1 has a biphasic  $\text{Ca}^{2+}$  response, and a two-site saturation fit yielded  $K_d$  values of  $0.81$  and  $60 \mu\text{M}$ . The two dissociation constants likely

result from the different affinities of the N- and C-terminal domains of CaM for  $\text{Ca}^{2+}$ , as observed for YC2.1 and YC4.3 (7). Fig. 1c shows that the cameleon fills a gap in  $\text{Ca}^{2+}$  sensitivity, falling directly between YC3.3 and YC4.3. Because the ER is expected to range from hundreds of micromolar (resting state) to low micromolar (depleted state), this sensor is ideally suited for imaging  $\text{Ca}^{2+}$  in the ER.

An important part of our design strategy was to develop a sensor that would not be perturbed by endogenous cellular proteins, such as WT CaM, because this has posed a problem for previous cameleons (21). Fig. 1d shows the FRET response of YC3.3 and the D1 in the presence of excess CaM. For YC3.3, the FRET response under saturating  $\text{Ca}^{2+}$  conditions decreases with the addition of increasing concentrations of CaM, because excess CaM in solution binds to the skMLCK portion of the cameleon, forming an intermolecular complex and preventing the sensor from registering a FRET increase. Importantly, in the charge reversal redesign (D1), addition of increasing CaM has almost no effect on the FRET response, indicating that this cameleon is not perturbed by large excesses of CaM.

Stopped-flow fluorescence measurements were undertaken to determine the binding kinetics of the cameleon. Rapid mixing of the cameleon protein with various concentrations of  $\text{Ca}^{2+}$  resulted in a rapid increase in FRET. The initial responses were fitted with a single exponential to obtain the observed rate ( $k_{\text{obs}}$ ). A fit to the data (Fig. 9, which is published as supporting information on the PNAS web site) yields  $k_{\text{on}}$  of  $3.6 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$ ,  $k_{\text{off}}$  of  $256 \text{ s}^{-1}$ , and  $K_d$  of  $69 \mu\text{M}$  for D1, where the observed  $K_d$  corresponds well to the results from the  $\text{Ca}^{2+}$  titration curve. Therefore, although this cameleon has a  $k_{\text{on}}$  similar to previous cameleons, it has a much faster  $k_{\text{off}}$  ( $250 \text{ s}^{-1}$  instead of  $10 \text{ s}^{-1}$ ), making it more appropriate to monitor rapidly changing  $\text{Ca}^{2+}$  dynamics.

Addition of the calreticulin signal sequence and a KDEL ER-retention tag led to effective and specific localization of the cameleon to the ER in mammalian cells (Fig. 2a). Fig. 2b and c show the dramatic improvement in  $[\text{Ca}^{2+}]_{\text{ER}}$  sensitivity of




**Fig. 3.** The effect of Bcl-2 overexpression on the  $[Ca^{2+}]_{ER}$  of MCF-7 cells. (a) Comparison of  $[Ca^{2+}]_{ER}$  in neo (red), Bcl-2 (blue), and Bcl-2 also overexpressing SERCA2b (green) cells. Each trace represents the average of more than five cells. Emission ratios were converted into  $[Ca^{2+}]$ , as described in *Materials and Methods*. Thapsigargin was added at time  $t = 0$  in the absence of external  $Ca^{2+}$  to prevent  $Ca^{2+}$  influx due to capacitative  $Ca^{2+}$  entry. (b) Leakage rate of  $Ca^{2+}$  from the ER as a function of  $[Ca^{2+}]_{ER}$  for Bcl-2 (blue), neo (red), and Bcl-2 overexpressing SERCA2b (green) cells.

D1ER over YC4.3ER, which was previously the best indicator for directly monitoring  $[Ca^{2+}]_{ER}$  at the single-cell level. Treatment of HeLa cells with pharmacological blockers such as thapsigargin, the irreversible sarcoplasmic reticulum  $Ca^{2+}$  ATPase (SERCA) inhibitor (Fig. 2*b*), or with agonists such as ATP, which results in  $IP_3$  production and activation of  $IP_3R$  (Fig. 2*c*), causes a depletion in  $[Ca^{2+}]_{ER}$  that is robust and readily visible by using D1ER. Additionally, upon ATP treatment, oscillations in  $[Ca^{2+}]_{ER}$  were directly observed with D1ER, whereas YC4.3ER was not sensitive enough to detect oscillations. The D1ER cameleon is not saturated under resting conditions, because treatment with 25  $\mu M$  digitonin and 5 mM  $Ca^{2+}$  led to a ratio increase (data not shown); thus, D1ER is poised to detect both increases and decreases in  $[Ca^{2+}]_{ER}$ .

Because CFP and citrine are spectrally distinct from the excitation and emission of fura-2, D1ER can be used in conjunction with this high-affinity  $Ca^{2+}$  indicator in the cytosol to simultaneously monitor both  $[Ca^{2+}]_{ER}$  and  $[Ca^{2+}]_{cyt}$  (Fig. 2*d* and *e*). As can be seen from Fig. 2*d*, in HeLa cells upon stimulation with 10  $\mu M$  ATP, there is a rise in  $[Ca^{2+}]_{cyt}$  followed by rapid oscillations. These oscillations are also observed in the ER, and, as evidenced by the data in Fig. 2*d* Inset, the  $[Ca^{2+}]_{ER}$  oscillations are directly anticorrelated with those in the cytosol. Dampening out of the oscillations coincides with a large decrease in  $[Ca^{2+}]_{ER}$ . At longer time points (Fig. 2*e*), once the  $[Ca^{2+}]_{cyt}$  has returned to essentially basal levels, lower-frequency sustained oscillations are observed in the cytosol, as described (24). We can now observe that these oscillations occur only when the ER starts to refill, presumably due to capacitative  $Ca^{2+}$  entry.


Given the dramatic improvement in our ability to directly measure  $[Ca^{2+}]_{ER}$  at the single-cell level with D1ER and the potential to monitor  $Ca^{2+}$  simultaneously in the ER and the cytosol, we turned our focus to the effect of the antiapoptotic Bcl-2 protein on  $[Ca^{2+}]_{ER}$  in MCF-7 breast cancer epithelial cell lines stably expressing either a control plasmid (neo) or Bcl-2 (25). As reported (25), overexpression of Bcl-2 in MCF-7 cells



**Fig. 4.** Comparison between ATP-induced  $Ca^{2+}$  oscillations in the cytosol and the ER for Bcl-2 and neo MCF-7 cells. The differences between Bcl-2 (a and c) and neo (b and d) cells are highlighted in c and d, which are on the same timescale and focus on the oscillations induced by ATP. The  $[Ca^{2+}]_{cyt}$  signal is in green, and the  $[Ca^{2+}]_{ER}$  signal is in blue for Bcl-2 and red for neo cells. The fura-2 excitation ratio (350/380 nm) and cameleon emission ratio (citrine/CFP) were converted to  $[Ca^{2+}]$ , as outlined in *Materials and Methods*.

protected against apoptosis induced by thapsigargin, staurosporine, and  $H_2O_2$ . Fig. 3*a* shows that in MCF-7 Bcl-2 cells,  $[Ca^{2+}]_{ER}$  is decreased by  $\approx 40\%$  [from an average of  $103 \pm 12 \mu M$  in neo cells ( $n = 45$  cells) to  $66 \pm 7 \mu M$  in Bcl-2 cells ( $n = 79$  cells)]. Overexpression of SERCA2b in the Bcl-2 cells restores  $[Ca^{2+}]_{ER}$  to control levels. Blocking SERCA pumps with thapsigargin causes depletion of  $[Ca^{2+}]_{ER}$  due to leak of  $Ca^{2+}$  across the ER membrane (Fig. 3*a*). Taking the derivative of the change in  $[Ca^{2+}]_{ER}$  upon thapsigargin treatment enables a comparison of the leakage rate for neo and Bcl-2 cells, and Bcl-2 cells overexpressing SERCA2b. As is evident from Fig. 3*b*, the leakage rate for Bcl-2 cells is greater than that for neo cells, indicating that under resting conditions, there is less  $[Ca^{2+}]_{ER}$  in Bcl-2 cells, because Bcl-2 increases the leak of  $Ca^{2+}$  out of the ER, as has been found in some other cell types (10, 11, 26). In cells overexpressing both Bcl-2 and SERCA2b, the leakage rate is still much greater than in control cells, indicating that the increase in resting  $[Ca^{2+}]_{ER}$  levels is due to increased pumping of  $Ca^{2+}$  into the ER rather than a decreased leak.

The molecular mechanism by which Bcl-2 alters  $Ca^{2+}$  homeostasis is still unclear. Bcl-2 may facilitate  $Ca^{2+}$  leakage from



**Fig. 5.** The effect of the green tea compound EGCG and a control compound EC on apoptosis and  $[Ca^{2+}]_{ER}$  of MCF-7 cells. (a) Cell viability (percent of the control), as determined by FACS analysis of propidium iodide and annexin V-stained neo (red) and Bcl-2 (blue) cells after 48-h treatment with nothing (control), EC, and increasing concentrations of EGCG. (b) Treatment of MCF-7 cells overexpressing Bcl-2 with EGCG and EC, as denoted by the green and blue boxes. The traces represent two different cells. (c) Treatment of MCF-7 neo cells with EGCG, as denoted by the green box. The traces represent two different cells. It should be noted that EC did not cause an increase in  $[Ca^{2+}]_{ER}$  in neo cells. (d) Bcl-2 cells treated with EGCG (green traces, each representing an individual cell) compared with neo (red) and Bcl-2 (blue) cells. At time  $t = 0$ , cells were treated with  $2 \mu\text{M}$  thapsigargin in the absence of external  $Ca^{2+}$  to determine the ER  $Ca^{2+}$  leakage rate. (e) Comparison of the leakage rates of  $Ca^{2+}$  from the ER for Bcl-2 (blue circles), neo (red diamonds), and Bcl-2 cells pretreated with EGCG (green, upside-down triangles and squares), showing the decreased leakage rate upon EGCG treatment.

the ER by forming a channel, either by homooligomerizing or by interacting with another member of the Bcl-2 family, because it has been reported that in planar lipid bilayers, Bcl-2 can form a cation selective channel at physiological pH (27). Alternatively, Bcl-2 could interact with and alter the function of an endogenous release channel or pore-forming protein in the ER. Indeed, in a recent study, Bcl-2 was found to interact with the IP<sub>3</sub>R in WEHI7.2 T cells by coimmunoprecipitation, and this interaction appeared to influence the amount of  $Ca^{2+}$  released from the ER and the open probability of the IP<sub>3</sub>R in lipid bilayers (16). If Bcl-2 interacts with the IP<sub>3</sub>R in MCF-7 cells, it would potentially alter  $Ca^{2+}$  oscillations triggered by ATP. Any differences between Bcl-2 and neo cells could be detected by simultaneously monitoring the ER and the cytosol.

Treatment of MCF-7 cells with ATP results in a large release of  $Ca^{2+}$  from the ER into the cytosol, followed by  $Ca^{2+}$  oscillations in both the ER and the cytosol upon refilling of the ER (Fig. 4). Bcl-2 cells required a larger dose of ATP than neo cells for oscillations to be induced (100 vs. 10  $\mu\text{M}$ ), suggesting that they are less sensitive to ATP. Fig. 4 *a* and *b* show responses of representative single Bcl-2 and neo cells. It is evident that less  $Ca^{2+}$  is released into the cytosol in Bcl-2 cells, both in the initial dump and in subsequent oscillations. The average initial decrease in  $[Ca^{2+}]_{ER}$  is  $51 \pm 4 \mu\text{M}$  for Bcl-2 cells and  $85 \pm 9 \mu\text{M}$  for neo cells, resulting in average cytosolic increases of  $890 \pm 370 \text{ nM}$  for Bcl-2 and  $1580 \pm 650 \text{ nM}$  for neo ( $n = 12$  for Bcl-2 cells and  $n = 11$  for neo cells). Importantly, in both Bcl-2 and neo cells,  $[Ca^{2+}]_{ER}$  decreases to  $\approx 5 \mu\text{M}$  after the initial release; therefore, the smaller  $Ca^{2+}$  release in Bcl-2 cells results from the lower  $Ca^{2+}$  level in the ER under resting conditions.

Aside from the intensity of the  $Ca^{2+}$  oscillations, the most significant difference between the Bcl-2 and neo cells is that the oscillations are typically more rapid and the duration of ER  $Ca^{2+}$  release is shorter in Bcl-2 overexpressing cells (Fig. 4 *c* and *d*).

As a consequence, Bcl-2 cells tend to oscillate at a higher frequency. Oscillation data from more individual cells are presented in Fig. 10, which is published as supporting information on the PNAS web site, demonstrating that, although there is some variability in the oscillations, the trends in oscillation frequency and duration are maintained.

Identifying novel inhibitors of Bcl-2 should provide insight into the role of Bcl-2 in endogenous signaling cascades leading to apoptosis, as well as lead to the potential development of therapeutic agents. Previously, Leone *et al.* (28) found that (-)EGCG, the most abundant component of green tea, could bind to the hydrophobic pocket of Bcl-2 *in vitro* and inhibit its interaction ( $K_i = 335 \text{ nM}$ ) with a peptide derived from the proapoptotic BH3-only protein Bad, whereas a compound lacking the gallate moiety, (-)EC, had no effect (see Fig. 11, which is published as supporting information on the PNAS web site, for structures). We treated both neo and Bcl-2 cells with EGCG and found that the compound could induce apoptosis in a dose-dependent manner, as determined by FACS analysis of propidium iodide- and annexin V-stained cells (Fig. 5*a*; details of assay given in *Supporting Text*). Importantly, EGCG was much more effective than other proapoptotic stimuli such as thapsigargin and  $H_2O_2$  in inducing apoptosis in Bcl-2 cells, indicating that it was capable of overcoming protection by Bcl-2. Given that green tea polyphenols have been shown to inhibit tumor formation and growth in certain models (29), this raises the possibility that this antitumor activity could be partially related to overcoming Bcl-2 protection of cancer cells.

To gain more insight into how EGCG overcomes Bcl-2 protection, we examined the effect of EGCG on  $[Ca^{2+}]_{ER}$ . As can be shown from Fig. 5*b*, treatment of Bcl-2 cells with  $150 \mu\text{M}$  EGCG results in an immediate increase in  $[Ca^{2+}]_{ER}$ , from an average of  $\approx 60$  to  $\approx 125 \mu\text{M}$ . This effect can be reversed upon washing out the inhibitor. On the contrary, the control com-

pound, EC, has no effect on  $[Ca^{2+}]_{ER}$ . Fig. 5c shows that in neo cells there is either no increase or a very modest increase in  $[Ca^{2+}]_{ER}$ , consistent with inhibition of the small amount of endogenous Bcl-2 in these cells. To determine the cause of the  $Ca^{2+}$  increase, we pretreated Bcl-2 cells with EGCG and then added thapsigargin (Fig. 5d) to monitor the leak rate. As can be seen from Fig. 5e, EGCG causes a dramatic decrease in the  $Ca^{2+}$  leak rate of Bcl-2 cells, such that the leak is the same or slower than that in neo cells. This suggests that the binding of EGCG to Bcl-2 prevents Bcl-2 from increasing the  $Ca^{2+}$  leak rate.

## Discussion

The improved ER-targeted cameleon poses a number of significant advantages over all previous methods for monitoring  $[Ca^{2+}]_{ER}$ . Compared with the ER-targeted aequorin probes, the cameleon signal is ratiometric, permits single-cell resolution, and is compatible with fura-2 in the same cell. Importantly, these characteristics enable the monitoring of  $Ca^{2+}$  oscillations within the ER. Additionally, the cameleon is not consumed by  $Ca^{2+}$  as aequorin is, and therefore the instantaneous signal depends only on  $Ca^{2+}$  and not the entire past history of  $Ca^{2+}$  in the cell (1). Because the cameleons and aequorins have different  $Ca^{2+}$  response curves, they should yield different mean  $[Ca^{2+}]$  whenever there is spatial heterogeneity in  $[Ca^{2+}]$ . The cameleons would tend to give slightly lower mean  $[Ca^{2+}]$  values than aequorins.

The improved ER-targeted cameleon enabled us to show that Bcl-2 plays an important role in governing  $[Ca^{2+}]_{ER}$  in breast cancer cells by altering the resting level of  $[Ca^{2+}]_{ER}$  as well as the amplitude, duration, and frequency of  $IP_3$ -mediated  $Ca^{2+}$  oscillations. Our study does not directly address whether Bcl-2 interacts with the  $IP_3R$  in MCF-7 cells, because  $Ca^{2+}$  oscillations are controlled by a host of factors, including the level of  $Ca^{2+}$  in the ER and capacitative calcium entry, both of which are altered in Bcl-2-overexpressing cells (9, 11, 26). However, our data show that, in addition to altering the resting level and leak rate of  $[Ca^{2+}]_{ER}$ , Bcl-2 overexpression can also modify fundamental  $Ca^{2+}$  signaling processes such as the nature of  $IP_3$ -induced  $Ca^{2+}$  oscillations. This raises the intriguing possibility that Bcl-2 may have an effect on signaling pathways that are downstream of  $Ca^{2+}$  oscillations and therefore has the potential to influence processes such as exocy-

tos, mitochondrial redox state, and differential gene transcription (30–32). This disruption of  $Ca^{2+}$  homeostasis by Bcl-2 may be essential in determining the susceptibility of breast cancer cells to different apoptotic stimuli, because Scorrano *et al.* (12) have found that certain apoptotic stimuli (lipid second messengers and oxidative stress, staurosporine, etoposide, and brefeldin-A) depend either critically or partially on ER  $Ca^{2+}$ .

We have also identified a small-molecule inhibitor (EGCG) of Bcl-2 that can overcome Bcl-2 protection and induce apoptosis in breast cancer cells. Interestingly, this inhibitor also reverses the effect of Bcl-2 on  $[Ca^{2+}]_{ER}$ . Because EGCG is known to bind in the hydrophobic groove and prevent the interaction of Bcl-2 with BH3-only peptides, this suggests that binding in this hydrophobic pocket inhibits the channel function of Bcl-2, either by preventing its interaction with other proteins or by causing a conformational change in Bcl-2 itself. Recently, Bassik *et al.* (33) found that phosphorylation within the unstructured loop region between the BH3 and BH4 domains of Bcl-2 inhibited the interaction between Bcl-2 and proapoptotic family members and caused an increase in  $Ca^{2+}$  leak rate (33). They also showed that the ability to bind BH3-only proteins, but not Bax or Bak, was critical for Bcl-2 to lower  $[Ca^{2+}]_{ER}$ . The combination of that study with the present one strongly suggests that Bcl-2 must interact with other proteins (either itself or other family members) through the hydrophobic groove to facilitate leakage of  $Ca^{2+}$  from the ER. It also implies that the interaction of other family members with Bcl-2 localized to the ER can have a dramatic effect on  $[Ca^{2+}]_{ER}$  and potentially apoptosis. We are currently examining whether a direct connection exists between blockage of the channel function, increase in  $[Ca^{2+}]_{ER}$ , and the increased apoptosis observed in the Bcl-2-overexpressing cells.

We thank L. Xu, S. R. Adams, L. Gross, P. Steinbach, and Q. Xiong for assistance and advice and Professor J. Adams for the use of his Applied Photophysics stopped-flow instrument. This work was supported by Ruth L. Kirschstein National Research Service Award Postdoctoral Fellowship F32-GM067488-01 (to A.E.P.), Department of Defense Postdoctoral Fellowship DAMD17-99-1-9096 (to C.J.), Department of Energy Grant DE-FG-01-ER63276 (to R.Y.T.), and National Institutes of Health Grants NS27177 (to R.Y.T.) and GM60554 (to J.C.R.).

- Alvarez, J. & Montero, M. (2002) *Cell Calcium* **32**, 251–260.
- Solovyova, N. & Verkhratsky, A. (2002) *J. Neurosci. Methods* **122**, 1–12.
- Sneyd, J., Tsaneva-Atanasova, K., Yule, D. I., Thompson, J. L. & Shuttleworth, T. J. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 1392–1396.
- Arnaudeau, S., Kelley, W. L., Walsh, J. V., Jr., & Demaurex, N. (2001) *J. Biol. Chem.* **276**, 29430–29439.
- Filippin, L., Magalhaes, P. J., Benedetto, G. D., Colella, M. & Pozzan, T. (2003) *J. Biol. Chem.* **278**, 39224–39234.
- Brini, M., Pinton, P., Pozzan, T. & Rizzuto, R. (1999) *Microsc. Res. Tech.* **46**, 380–389.
- Miyawaki, A., Llopis, J., Heim, R., McCaffery, J. M., Adams, J. A., Ikura, M. & Tsien, R. Y. (1997) *Nature* **388**, 882–887.
- Cory, S. & Adams, J. M. (2002) *Nat. Rev. Cancer* **2**, 647–656.
- Lam, M., DUBYAK, G., Chen, L., Nunez, G., Miesfeld, R. & Distelhorst, C. W. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 6569–6573.
- Foyouzi-Youssefi, R., Arnaudeau, S., Borner, C., Kelly, W. L., Tschopp, J., Lew, D. P., Demaurex, N. & Krause, K.-H. (2000) *Proc. Natl. Acad. Sci. USA* **97**, 5723–5728.
- Pinton, P., Ferrari, D., Magalhaes, P., Schulze-Osthoff, K., Virgilio, F. D., Pozzan, T. & Rizzuto, R. (2000) *J. Cell Biol.* **148**, 857–862.
- Scorrano, L., Oakes, S. A., Opferman, J. T., Chang, E. H., Sorcinelli, M. D., Pozzan, T. & Korsmeyer, S. J. (2003) *Science* **300**, 135–139.
- Baffy, G., Miyashita, T., Williamson, J. R. & Reed, J. C. (1993) *J. Biol. Chem.* **268**, 6511–6519.
- Pinton, P., Ferrari, D., Rapizzi, E., Virgilio, F. D., Pozzan, T. & Rizzuto, R. (2001) *EMBO J.* **20**, 2690–2701.
- Distelhorst, C. W. & Shore, G. C. (2004) *Oncogene* **23**, 2875–2880.
- Chen, R., Valencia, I., Zhong, F., McColl, K. S., Roderick, H. L., Bootman, M. D., Berridge, M. J., Conway, S. J., Holmes, A. B., Mignery, G. A., *et al.* (2004) *J. Cell Biol.* **166**, 193–203.
- Griesbeck, O., Baird, G. S., Campbell, R. E., Zacharias, D. A. & Tsien, R. Y. (2001) *J. Biol. Chem.* **276**, 29188–29194.
- Tsien, R. Y. & Pozzan, T. (1989) *Methods Enzymol.* **172**, 230–263.
- Tsien, R. Y. & Rink, T. J. (1983) in *Current Methods in Cellular Neurobiology*, ed. Barber, J. L. (Wiley, New York), Vol. 3, pp. 249–312.
- Poenie, M. (1990) *Cell Calcium* **11**, 85–91.
- Hasan, M. T., Friedrich, R. W., Larkum, M. E., Euler, T., Both, M., Duebel, J., Waters, J., Giese, G., Bujard, H., Griesbeck, O., *et al.* (2004) *Pub. Lib. Sci.* **2**, 763–775.
- Heim, N. & Griesbeck, O. (2004) *J. Biol. Chem.* **279**, 14280–14286.
- Ikura, M., Clore, G. M., Gronenborn, A. M., Zhu, G., Klee, C. B. & Bax, A. (1992) *Science* **256**, 632–638.
- Bootman, M. D., Cheek, T. R., Moreton, R. M., Bennett, D. L. & Berridge, M. J. (1994) *J. Biol. Chem.* **269**, 24783–24791.
- Froesch, B. A., Aime-Sempe, C., Leber, B., Andrews, D. & Reed, J. C. (1999) *J. Biol. Chem.* **274**, 6469–6475.
- Abeele, F. V., Skryma, R., Shuba, Y., Coppenolle, F. V., Slomianny, C., Roudbaraki, M., Mauroy, B., Wuytack, F. & Prevarskaya, N. (2002) *Cancer Cell* **1**, 169–179.
- Schendel, S. L., Xie, Z., Montall, M. O., Matsuyama, S., Montal, M. & Reed, J. C. (1997) *Proc. Natl. Acad. Sci. USA* **94**, 5113–5118.
- Leone, M., Zhai, D., Sareth, S., Kitada, S., Reed, J. C. & Pellecchia, M. (2003) *Cancer Res.* **63**, 8118–8121.
- Yang, C. S. & Wang, Z. Y. (1993) *J. Natl. Cancer Inst.* **85**, 1038–1049.
- Berridge, M. J., Bootman, M. D. & Roderick, H. L. (2003) *Nat. Mol. Cell Biol.* **4**, 517–529.
- Berridge, M. J., Lipp, P. & Bootman, M. D. (2000) *Nat. Mol. Cell Biol.* **1**, 11–21.
- Blaustein, M. P. & Golovina, V. A. (2001) *Trends Neurosci.* **24**, 602–608.
- Bassik, M. C., Scorrano, L., Oakes, S. A., Pozzan, T. & Korsmeyer, S. J. (2004) *EMBO J.* **23**, 1207–1216.